WALAILAK JOURNAL

A Fractional Model of Bloch Equation in Nuclear Magnetic Resonance and its Analytic Approximate Solution

Sunil KUMAR^{1,*}, Naeem FARAZ² and Khosro SAYEVAND³

¹Department of Mathematics, National Institute of Technology, Jamshedpur, Jharkhand, India ²Modern Textile Institute, Donghua University, Yan'an Xilu Road, Shanghai, China ³Department of Mathematics, Faculty of Basic Sciences, Malayer University, Malayer, Iran

(*Corresponding author's e-mail: skiitbhu28@gmail.com, skumar.math@nitjsr.ac.in)

Abstract

The purpose of this paper is to employ an analytical approach to the time fractional Bloch Nuclear Magnetic Resonance (NMR) flow equations. A comparative study of the numerical solutions and the well-known analytical solutions are discussed. Absolute error has been calculated to show the accuracy of the applied method. The numerical solutions show that only a few iterations are needed to obtain accurate approximate solutions. The fractional derivatives are described in the Caputo sense. Numerical results are presented graphically.

Keywords: Bloch equation, Caputo derivative, analytical solution, homotopy perturbation method

Introduction

In physics and chemistry, specifically in nuclear magnetic resonance (NMR), magnetic resonance imaging (MRI), and electron spin resonance (ESR), the Bloch equations are a set of macroscopic equations that are used to calculate the nuclear magnetization $M = (M_x, M_y, M_z)$ in the laboratory frame (x,y,z). T_1 and T_2 are known respectively as the spin lattice and sin-spin relaxation times which measure the interaction of the nuclei with their surroundings molecular environment and those between close nuclei. The MRI is a powerful tool for obtaining spatially localized information from the NMR of atoms within a sample. Sometimes they are called the equations of motion of nuclear magnetization. These equations were introduced by Felix Bloch [1]. In 1956, Torrey modified the Bloch equations by incorporating a diffusion term by Torrey [2]. The dynamics of an ensemble of spins without mutual couplings is usually well described by the Bloch equations [3,4], which can be viewed as mathematical descriptions of precession of the macroscopic magnetization vector around a (possibly time-dependent) magnetic field. Recently, Petras [5] and Bhalekar *et al.* [6] solved the fractional Bloch equation. The Bloch equations can be expressed in the following form:

$\left[\frac{d}{d}\right]$	$\frac{M_x(t)}{dt} =$	$=\omega_0 M_y(t)-\frac{M_x(t)}{T_2},$
$\left \frac{d}{d}\right $	$\frac{dM_{y}(t)}{dt} =$	$=\omega_0 M_x(t) - \frac{M_y(t)}{T_2},$
$\left \frac{d}{d}\right $	$\frac{dM_z(t)}{dt} =$	$=\frac{M_0-M_z(t)}{T_1},$

(1)

with initial conditions $M_x(0) = 0$, $M_y(0) = 100$ and $M_z(0) = 0$, where $\omega_0 = \gamma B_0$ and $\omega_0 = 2\pi f_0$ (e.g. gyromagnetic ratio $\gamma/2\pi = f_0/B_0 = 42.57 MH_z/T$ (for water protons) and M_0 is the equilibrium magnetization. The complete set of analytic solutions of the system of Eq. (1) is given as [6];

$$\begin{cases}
M_{x}(t) = e^{-t/T_{2}} \left(M_{x}(0) \cos \omega_{0} t + M_{y}(0) \sin \omega_{0} t \right), \\
M_{y}(t) = e^{-t/T_{2}} \left(M_{y}(0) \cos \omega_{0} t - M_{x}(0) \sin \omega_{0} t \right), \\
M_{z}(t) = M_{z}(0) e^{-t/T_{1}} + M_{0}(1 - e^{-t/T_{1}}).
\end{cases}$$
(2)

Recently, many experts have paid great attention to construct the solutions of the Bloch equations by different methods from exact solutions to the Bloch equations by Bain *et al.* [7], analytical solution of the time dependent Bloch NMR by Awojoyogber [8], a solution to Bloch NMR flow equations by Awojoyogber and Boubaker [9], numerical solutions to the time dependent Bloch equations by Murase and Tanki [10], fast approximate solution of Bloch equation by Balac and Chupin [11], some Solutions of the Bloch equations by Leyte [12], solution of the Bloch equation by Hoult [13], approximate solutions of the Bloch equations by Yan *et al.* [14], exact solutions of the Bloch equations by Sivers [16], and a Near-Resonance Solution to the Bloch equations by Xu and Chan [17].

Mathematical modeling of many physical systems leads to linear and nonlinear fractional differential equations in various fields of physics and engineering. The use of fractional differentiation for the mathematical modeling of real world physical problems has been widespread in recent years, e.g. modeling of earthquakes, fluid dynamic traffic models with fractional derivatives, measurement of viscoelastic material properties, etc. The book by Oldham and Spanier [18] has played a key role in the development of the subject. Some fundamental results related to solving fractional differential equations may be found in books [19-21].

The main aim of this article presents a mathematical model of Bloch equations with fractional time derivative α, β, γ ($0 < \alpha, \beta, \gamma \le 1$) in the form of a rapidly convergent series with easily computable components. This method was first proposed and applied by He [22-25] and was successfully applied to solve many problems related to fractional calculus [26-31].

Basic Idea of Homotopy Perturbation Method (HPM)

To illustrate the basic ideas of the HPM for fractional differential equations, the following problem is considered:

$$D_{*t}^{a}\xi(x,t) = v(x,t) - L\xi(x,t) - N\xi(x,t), m - 1 < \alpha < m, m \in N, t \ge 0, x \in \mathbb{R}^{n},$$
(3)

subject to the initial and boundary conditions;

$$\xi^{(i)}(0,0) = c_i, \ B\left(\xi, \frac{\partial\xi}{\partial x_j}, \frac{\partial\xi}{\partial t}\right) = 0, i = 0, 1, \dots, m-1, \ j = 1, 2, \dots n,$$
(4)

where L is a linear operator, while N is a nonlinear operator, ν is a known analytical function and $D_{*_t}^{\alpha}$ denotes the fractional derivative in the Caputo sense. ξ is assumed to be a causal function of time, i.e.,

vanishing for t < 0. Also $\xi^{(i)}(x,t)$ is the *ith* derivative of $\xi, c_i, i = 0, 1, ..., m-1$ are the specified initial conditions, and *B* is a boundary operator.

In view of He's homotopy perturbation technique, the following simple homotopy can be constructed.

$$(1-p)D_{*_{t}}^{\alpha}\xi(x,t) + p\left(D_{*_{t}}^{\alpha}\xi(x,t) + L\xi(x,t) + N\xi(x,t) - \nu(x,t)\right) = 0, \ p \in [0,1],$$
or
$$(5)$$

$$D_{*t}^{\alpha}\xi(x,t) + p(L\xi(x,t) + N\xi(x,t) - \nu(x,t)) = 0, p \in [0,1].$$
(6)

The homotopy parameter p always changes from zero to unity. In case p = 0 Eq. (5) and Eq. (6) becomes;

$$D^{\alpha}_{*t}\xi(x,t) = 0, \tag{7}$$

where p = 1, Eq. (5) and Eq. (6) turns out to be the original fractional differential equation. In view of the homotopy perturbation method, the homotopy parameter p is used to expand the solution in the following form.

$$\xi(x,t) = \xi_0(x,t) + p\xi_1(x,t) + p^2\xi_2(x,t) + p^3\xi_3(x,t) + \dots$$
(8)

For a nonlinear problem, set $N\xi(x,t) = S(x,t)$ Substituting Eq. (8) into Eq. (6) or Eq. (5) and equating the terms with identical power of p, a series of equations of the form is obtained.

$$p^{0}: D_{*_{t}}^{\alpha}\xi_{0}(x,t) = 0,$$

$$p^{1}: D_{*_{t}}^{\alpha}\xi_{1}(x,t) = -L\xi_{0}(x,t) - S_{0}(\xi_{0}(x,t)) + v(x,t),$$

$$p^{2}: D_{*_{t}}^{\alpha}\xi_{2}(x,t) = -L\xi_{1}(x,t) - S_{1}(\xi_{0}(x,t),\xi_{1}(x,t)),$$

$$p^{3}: D_{*_{t}}^{\alpha}\xi_{3}(x,t) = -L\xi_{2}(x,t) - S_{2}(\xi_{0}(x,t),\xi_{1}(x,t),\xi_{2}(x,t)),$$
(9)

where the functions S_0, S_1, S_2, \dots satisfy the following equation.

$$S(\xi_0(x,t) + p\xi_1(x,t) + p^2\xi_2(x,t) + p^3\xi_3(x,t) + ...) = S_0(\xi_0(x,t)) + pS_1(\xi_0(x,t),\xi_1(x,t)) + p^2S_2(\xi_0(x,t),\xi_1(x,t),\xi_2(x,t)) + ...$$
(10)

Applying the inverse operator on both sides of the equation and considering the initial and boundary conditions, the terms of the series solution can be given by;

$$\begin{cases} \xi_{0}(x,t) = \sum_{i=0}^{n-1} c_{i} \frac{t^{i}}{i!}, \\ \xi_{1}(x,t) = -I_{t}^{\alpha} (L\xi_{0}(x,t)) - I_{t}^{\alpha} S_{0}(\xi_{0}(x,t)) + I_{t}^{\alpha} v(x,t), \\ \xi_{j}(x,t) = -I_{t}^{\alpha} (L\xi_{j-1}(x,t)) - I_{t}^{\alpha} S_{j-1}(\xi_{0}(x,t),\xi_{1}(x,t),...,\xi_{j-1}(x,t)), j = 2,3,... \end{cases}$$

$$(11)$$

Hence, an accurate approximate in the following form is obtained.

$$\xi(x,t) = \sum_{i=0}^{\infty} \xi_i(x,t) \,.$$
(12)

Basic definitions of fractional calculus

In this section, some basic definitions and properties of fractional calculus theory are given which are used in this paper;

Definition2.1. A real function f(t), t > 0 is said to be in the space C_{μ} , $\mu \in R$ if there exists a real number $p > \mu$, such that $f(t) = t^p f_1(t)$ where $f_1(t) \in C(0, \infty)$ and it is said to be in the space C_n if and only if $f^{(n)} \in C_{\mu}$, $n \in N$.

Definition2.2. The Riemann-Liouville fractional integral (J_t^{α}) operator of order α of a function $f \in C_{\mu}, \mu \geq -1$ is defined as;

$$\begin{cases} J_{t}^{\alpha} f(t) = \frac{1}{\Gamma(\alpha)} \int_{0}^{t} (t-\tau)^{\alpha-1} f(\tau) d\tau, \ (\alpha > 0, \ t > 0), \\ J_{t}^{\alpha} f(t) = f(t). \end{cases}$$
(13)

where $\Gamma(.)$ is the well-known Gamma function. Some of the properties of the operator (j_t^{α}) , can be found in [18-21]; only the following are considered. For $f \in C_{\mu}$, $\mu \ge -1$, α , $\beta \ge 0$ and $\gamma \ge -1$;

(1) $J^{\alpha}_{t}J^{\beta}_{t}f(t) = J^{\alpha+\beta}_{t}f(t),$

(2)
$$\left(J_t^{\alpha}J_t^{\beta}\right)f(t) = \left(J_t^{\beta}J_t^{\alpha}\right)f(t),$$

(3) $J_t^{\alpha} t^{\gamma} = \frac{\Gamma(\gamma+1)}{\Gamma(\gamma+\alpha+1)} t^{\gamma+\alpha}.$

The Riemann-Liouville derivative has certain disadvantages when trying to model real world phenomena with fractional differential equations.

Definition 2.3. The fractional derivative D_t^{α} of f(t) in the Caputo sense defined as;

$$D_{t}^{\alpha}f(t) = \frac{1}{\Gamma(m-\alpha)} \int_{0}^{t} \frac{f^{(m)}(\tau)}{(t-\tau)^{\alpha+1-m}} d\tau,$$
(14)

where $m-1 < \alpha \leq m, m \in N, t > 0, f \in C_{-1}^m$.

Walailak J Sci & Tech 2014; 11(4)

The following are 2 basic properties of the Caputo's fractional derivative;

Lemma2.1. If $m-1 < \alpha \le m$, $m \in N$ and $f \in C_u^n$, $\mu \ge -1$, then;

$$(D_{t}^{\alpha}J_{t}^{\alpha})f(t) = f(t),$$

$$(J_{t}^{\alpha}D_{t}^{\alpha})f(t) = f(t) - \sum_{i=0}^{m-1} f^{i}(0^{+})\frac{t^{i}}{i!},$$
(15)

Definition2.6. The Mittag-Leffler function $E_{\alpha}(z)$ with $\alpha > 0$ is defined by the following series representation, valid in the whole complex plane [32];

$$E_{\alpha}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha \ n+1)}, \qquad \alpha > 0, \quad z \in C.$$
(16)

Solution of the given problem by HPM

The fractional-order Bloch equations are considered where integer order derivatives are replaced by fractional order as;

$$\begin{cases} D_{t}^{\alpha}M_{x}(t) = \omega_{0}M_{y}(t) - \frac{M_{x}(t)}{T_{2}}, & 0 < \alpha \leq 1, \\ D_{t}^{\beta}M_{y}(t) = \omega_{0}M_{x}(t) - \frac{M_{y}(t)}{T_{2}}, & 0 < \beta \leq 1, \\ D_{t}^{\gamma}M_{z}(t) = \frac{M_{0} - M_{z}(t)}{T_{1}}, & 0 < \gamma \leq 1, \end{cases}$$
(17)

where α , β and γ are the derivative orders. The total order of the system is (α, β, γ) . Here, all parameters ω_0 , T_1 and T_2 have units of $(s)^{-q}$ to maintain a consistent set of units for the magnetization.

The following homotopy is constructed.

$$\begin{cases} D_t^{\alpha} M_x(t) = p \left(\omega_0 M_y(t) - \frac{M_x(t)}{T_2} \right), & 0 < \alpha \le 1, \\ D_t^{\beta} M_y(t) = p \left(-\omega_0 M_x(t) - \frac{M_y(t)}{T_2} \right), & 0 < \beta \le 1, \\ D_t^{\beta} M_z(t) = p \left(\frac{M_0 - M_z(t)}{T_1} \right), & 0 < \gamma \le 1, \end{cases}$$

$$(18)$$

with appropriate initial conditions $M_x(0) = 0$, $M_y(0) = 100$ and $M_z(0) = 0$.

Assuming the solution of the fractional-order Bloch equations (18) to be in the following form;

$$M_{x}(t) = \lim_{N \to \infty} \sum_{i=0}^{N} p^{i} M_{x,i}(t), \ M_{y}(t) = \lim_{N \to \infty} \sum_{i=0}^{N} p^{i} M_{y,i}(t), \ M_{z}(t) = \lim_{N \to \infty} \sum_{i=0}^{N} p^{i} M_{z,i}(t).$$
(19)

Walailak J Sci & Tech 2014; 11(4)

277

Sunil KUMAR *et al*.

Substituting Eq. (19) into Eq. (18) and collecting terms of the same power of p, the following set of differential equations is obtained.

$$p^{0}: \begin{cases} D_{t}^{\alpha}M_{x,0}(t) = 0, \\ D_{t}^{\beta}M_{y,0}(t) = 0, \\ D_{t}^{\gamma}M_{z,0}(t) = 0, \end{cases}$$
(20)

$$p^{1}: \begin{pmatrix} D_{t}^{\alpha}M_{x,1}(t) = \omega_{0}M_{y,0}(t) - \frac{M_{x,0}(t)}{T_{2}}, \\ D_{t}^{\beta}M_{y,1}(t) = -\omega_{0}M_{x,0}(t) - \frac{M_{y,0}(t)}{T_{2}}, \\ D_{t}^{\gamma}M_{z,1}(t) = \frac{M_{0} - M_{x,0}(t)}{T_{1}}, \end{pmatrix} p^{n}: \begin{cases} D_{t}^{\alpha}M_{x,n}(t) = \omega_{0}M_{y,n-1}(t) - \frac{M_{x,n-1}(t)}{T_{2}}, \\ D_{t}^{\beta}M_{y,n}(t) = -\omega_{0}M_{x,n-1}(t) - \frac{M_{y,n-1}(t)}{T_{2}}, \\ D_{t}^{\gamma}M_{z,n}(t) = -M_{z,n-1}, \end{cases}$$
(21)

Consequently, the above system of nonlinear equations can be easily solved by applying the operator J_t^{α} to (20) - (21), giving the various components $M_{xn}(t)$, $M_{yn}(t)$ and $M_{zn}(t)$, thus enabling the series solution to be entirely determined. The first few components of the homotopy perturbation solution for Eq. (19) are derived as follows.

$$M_{x0}(t) = M_x(0) = 0,$$
 $M_{y0}(t) = M_y(0) = 100,$ $M_{z0}(t) = M_z(0) = 0,$ (22)

$$M_{x1}(t) = \frac{100\omega_0 t^{\alpha}}{\Gamma(\alpha+1)}, \qquad M_{y1}(t) = -\frac{100}{T_2} \frac{t^{\beta}}{\Gamma(\beta+1)}, \qquad M_{z1}(t) = \frac{M_0}{T_1} \frac{t^{\gamma}}{\Gamma(\gamma+1)}, \quad (23)$$

$$M_{x2}(t) = -\frac{100\omega_0}{T_2} \left(\frac{t^{2\alpha}}{\Gamma(2\alpha+1)} + \frac{t^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} \right), M_{y2}(t) = 100 \left(\frac{t^{2\beta}}{T_2^2 \Gamma(2\beta+1)} - \frac{\omega_0^2 t^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} \right), M_{z2}(t) = -\frac{M_0}{T_1} \frac{t^{2\gamma}}{\Gamma(2\gamma+1)}, (24)$$

$$M_{x3}(t) = \frac{100\,\omega_0}{T_2^2} \left(\frac{t^{3\alpha}}{\Gamma(3\alpha+1)} + \frac{t^{\alpha+2\beta}}{\Gamma(\alpha+2\beta+1)} + \frac{t^{2\alpha+\beta}}{\Gamma(2\alpha+\beta+1)} \right) - \frac{100\,\omega_0^3 t^{2\alpha+\beta}}{\Gamma(2\alpha+\beta+1)},$$

$$M_{\gamma_3}(t) = \frac{100\,\omega_0^2}{T_2} \left(\frac{2t^{\alpha+2\beta}}{\Gamma(\alpha+2\beta+1)} + \frac{t^{2\alpha+\beta}}{\Gamma(2\alpha+\beta+1)} \right) - \frac{100}{T_2^3} \frac{t^{3\beta}}{\Gamma(3\beta+1)}, \quad M_{z3}(t) = \frac{M_0}{T_1} \frac{t^{3\gamma}}{\Gamma(3\gamma+1)}, \tag{25}$$

and so on. In this manner, the rest of components of the homotopy perturbation solution can be obtained. The approximate solutions $\tilde{M}_x(t)$, $\tilde{M}_y(t)$ and \tilde{M}_z , by truncating the respective solutions series at level N = 14, are given by;

$$\widetilde{M}_{x}(t) = \sum_{n=0}^{14} M_{x,n}(t), \ \widetilde{M}_{y}(t) = \sum_{n=0}^{14} M_{y,n}(t), \ \widetilde{M}_{z}(t) = \sum_{n=0}^{14} M_{z,n}(t),$$
(26)

hence,

278

$$\begin{cases} \tilde{M}_{x}(t) = 100 \ \omega_{0} \bigg(\frac{t^{\alpha}}{\Gamma(\alpha+1)} - \frac{t^{2\alpha}}{T_{2}\Gamma(2\alpha+1)} - \frac{t^{\alpha+\beta}}{T_{2}\Gamma(\alpha+\beta+1)} + \frac{t^{3\alpha}}{T_{2}^{2}\Gamma(3\alpha+1)} + \frac{t^{2\alpha+\beta}}{T_{2}^{2}\Gamma(2\alpha+\beta+1)} - \frac{\omega_{0}^{2}t^{2\alpha+\beta}}{\Gamma(2\alpha+\beta+1)} + \frac{t^{\alpha+2\beta}}{T_{2}^{2}\Gamma(\alpha+2\beta+1)} + \dots \bigg) \\ \tilde{M}_{y}(t) = 100 \ E_{\beta} \bigg(-\frac{t^{\beta}}{T_{2}} \bigg) - \frac{100 \ \omega_{0}^{2}t^{\alpha+\beta}}{\Gamma(\alpha+\beta+1)} + \frac{100 \ \omega_{0}^{2}}{T_{2}} \bigg(\frac{2t^{\alpha+\beta}}{\Gamma(\alpha+2\beta+1)} + \frac{t^{2\alpha+\beta}}{\Gamma(2\alpha+\beta+1)} \bigg) + \dots \end{aligned}$$

$$(27)$$

$$\tilde{M}_{z}(t) = \frac{M_{0}}{T_{1}} \bigg(1 - E_{y}(-t^{\gamma}) \bigg)$$

The series solution converges very rapidly. The rapid convergence means only a few terms are required to get the approximate solutions.



Figure 1 The Exact M_x (solid line), M_y (solid line) and the approximate solution \widetilde{M}_x (Red dotted line) and \widetilde{M}_x (Blue dotted) at $\omega = 1$ and $T_2 = 20$ $(ms)^q$.



Figure 2 The Exact M_z (solid line) and the approximate solution \tilde{M}_z (Red dotted line) at $\omega = 1$ and $T_1 = 1$ (*ms*)^{*q*}.

Walailak J Sci & Tech 2014; 11(4)

The observations are depicted through **Figures 1 - 8**. **Figure 1 - 2** shows the comparison between the exact solutions and approximate solutions (by HPM) for the standard Bloch equation, i.e. for $\alpha = \beta = \gamma = 1$. It can be seen from **Figures 1 - 2** that the approximate solutions $\tilde{M}_x(t), \tilde{M}_y(t), \tilde{M}_z(t)$ obtained by the present method are nearly identical to the exact solutions $M_x(t), M_y(t), M_y(t)$ with high accuracy. All graphical result have parameters $\omega = 1$, $T_1 = 1(s)^q$ and $T_2 = 20$ (ms)^q.

Numerical result and discussion

In this section, the error analysis between the exact solutions and approximate solutions are depicted through **Figures 3** – **5**, and **Figures 6** - **8** shows the approximate solutions for different fractional Brownian motions and standard motions. During numerical computation, only 14 terms of the series solution are considered. The accuracy of the result can be improved by introducing more terms of the approximate solutions.

The simplicity and accuracy of the proposed method is illustrated by computing the absolute errors $e_{M_x}(t) = |M_x(t) - \tilde{M}_x(t)|$, $e_{M_y}(t) = |M_y(t) - \tilde{M}_y(t)|$, and $e_{M_z}(t) = |M_z(t) - \tilde{M}_z(t)|$, where $M_x(t)$, $M_y(t) M_z(t)$ are the exact solutions and $\tilde{M}_x(t)$, $\tilde{M}_y(t)$, $\tilde{M}_z(t)$ are the approximate solutions of (3) obtained by truncating the respective solutions series (21) at level N = 14. Figures 3 - 5 represent absolute errors between the exact solutions $M_x(t)$, $M_y(t)$, $M_y(t)$, $M_y(t)$, $M_y(t)$, $M_y(t)$, and the approximate solutions $\tilde{M}_x(t)$, $\tilde{M}_y(t)$, $\tilde{M}_z(t)$ respectively, which show high accuracy of the approximate solutions. Figures 3 - 5 show that the series solution (by HPM) converges to the exact solution very rapidly.



Figure 3 Absolute error $e_x = |M_x(t) - \tilde{M}_x(t)|$ at $\omega = 1$ and $T_2 = 20$ $(ms)^q$.



Figure 4 Absolute error $e_y = |M_y(t) - \widetilde{M}_y(t)|$ at $\omega = 1$ and $T_2 = 20 \ (ms)^q$.



Figure 5 Absolute error $e_z = |M_z(t) - \tilde{M}_z(t)|$ at $\omega = 1$ and $T_1 = 1$ $(ms)^q$.



Figure 6 Approximate solution $\tilde{M}_{x}(t)$ for different values of α at $\omega = 1$ and $T_2 = 20 \ (ms)^q$.



Figure 7 Approximate solution $\tilde{M}_{y(t)}$ for different values of β at $\omega = 1$ and $T_2 = 20$ $(ms)^q$.



Figure 8 Approximate solution $\tilde{M}_{r}(t)$ for different values of γ at $\omega = 1$ and $T_{1} = 1$ $(ms)^{q}$.

Figures 6 - 8 show the evolution of results for the approximate solutions $\tilde{M}_x(t)$, $\tilde{M}_y(t)$ and $\tilde{M}_z(t)$ of Eq. (3) obtained for different values α, β and γ respectively, by using the HPM. It is seen from Figures 6 and 8 that the approximate solutions $\tilde{M}_x(t)$ and $\tilde{M}_z(t)$ increase with the increases in t for different value of $\alpha = \beta = 0.7$, 0.8, 0.9, and also for the standard Bloch equation i.e. for $\alpha = \beta = 1$. It is also seen from Figure 7 that the approximate solution $\tilde{M}_y(t)$ decreases with the increases in t for different values of $\gamma = 0.7$, 0.8, 0.9, 1. It is to be noted that only 14 terms of the homotopy perturbation series were used in evaluating the approximate solutions in all figures.

Conclusions

The present case studied the effect of different orders of fractional time for the Bloch NMR flow equations, which are a set of macroscopic equations that are used for modeling of nuclear magnetization as a function of time by homotopy perturbation method. The following observations have been made.

1) $\tilde{M}_{x}(t)$ and $\tilde{M}_{z}(t)$ increase with the increases in t for different values of $\alpha = \beta = 0.7, 0.8, 0.9$.

2) For the standard Bloch equation i.e. for $\alpha = \beta = 1$. $\tilde{M}_x(t)$ and $\tilde{M}_z(t)$ increase with the increases in t.

3) Absolute errors between the exact solutions $M_x(t)$, $M_y(t)$, $M_y(t)$, and approximate solutions $\tilde{M}_x(t)$, $\tilde{M}_y(t)$, $\tilde{M}_z(t)$ respectively show high accuracy of the approximate solutions.

Acknowledgement

The author is very grateful to the anonymous referees for carefully reading the paper and for their constructive comments and suggestions which have improved the paper.

References

- [1] F Bloch. Nuclear induction. Phys. Rev. 1946; 70, 460-74.
- [2] HC Torrey. Bloch equations with diffusion terms. Phys. Rev. 1956; 104, 563-5.
- [3] J Jeener. Dynamical effects of the dipolar field in homogeneities in high-resolution NMR: spectral clustering and instabilities. *Phys. Rev. Lett.* 1999; **82**, 1772-5.
- [4] DE Rourke, L Khodarinova and AA Karabanov. Two-level systems with relaxation. *Phys. Rev. Lett.* 2004; **92**, 1-4.
- [5] I Petras. Modelling and numerical analysis of fractional order Bloch equations. *Comput. Math. Appl.* 2011; **61**, 341-56.
- [6] S Bhalekar, GV Daftardar, D Baleanu and R Magin. Fractional Bloch equation with delay. *Comput. Math. Appl.* 2011; **61**, 1355-65.
- [7] AD Bain, CK An and Z Nie. Exact solution to the Bloch equations and application to the Hahn echo. *J. Magn. Res.* 2010; **206**, 227-40.
- [8] OB Awojoyogber. Analytical solution of the time dependent Bloch NMR, flow equations: a translational mechanical analysis. *Phys. A Stat. Mech. Appl.* 2004; **339**, 437-60.
- [9] OB Awojoyogber and K Boubaker. A solution to Bloch NMR flow equations for the Analysis of hemodynamics functions of blood flow system using m-Boubaker polynomials. *Curr. Appl. Phys.* 2009; 2, 278-83.
- [10] K Murase and N Tanki. Numerical solutions to the time dependent Bloch equations revisited. *Magn. Res. Imag.* 2011; **29**, 126-31.
- [11] S Balac and L Chupin. Fast approximate solution of Bloch equation for simulation of RF artifacts in Magnetic Resonance Imaging. *Math. Comput. Model.* 2008; **48**, 1901-13.
- [12] JC Leyte. Some solutions of the Bloch equations. Chem. Phys. Lett. 1990; 165, 213-20.
- [13] DI Hoult. The solution of the Bloch equation in the presences of a varying B1 field An approach to selective pulse analysis. *J. Magn. Res.* 1979; **35**, 69-86.
- [14] H Yan, B Chen and JC Gore. Approximate solutions of the Bloch equations for selective excitation. *J. Magn. Res.* 1987; **75**, 83-95.
- [15] J Schotland and JS Leigh. Exact solution of the Bloch equations with n-site chemical exchange. J. Magn. Res. 1983; 51, 48-55.
- [16] EA Sivers. Approximate solution to the Bloch equation with symmetric RF pulses and flip angles less than TT/2. *J. Magn. Res.* 1986; **69**, 28-40.
- [17] Z Xu and AK Chan. A Near-Resonance Solution to the Bloch equations and its application to RF pulse design. J. Magn. Res. 1999; 138, 225-31.
- [18] KB Oldham and J Spanier. The Fractional Calculus. Academic Press, New York, 1974.
- [19] KS Miller and B Ross. An Introduction to the Fractional Calculus and Fractional Differential *Equations*. John Willey and Sons, New York, 2003.
- [20] I Podlubay. Fractional Differential Equations Calculus. Academic, Press, New York.
- [21] AA Kilbas, HM Srivastava and JJ Trujillo. *Theory and Application of Fractional Differential Equations*. Elsevier, Amsterdam, 2006.
- [22] JH He. Homotopy perturbation technique. Comput. Meth. Appl. Mech. Eng. 1999; 178, 257-62.
- [23] JH He. A coupling of method of homotopy technique and perturbation technique for nonlinear problems. *Int. J. Nonlinear Mech.* 2000; **35**, 37-43.
- [24] JH He. Comparison of homotopy perturbation method and homotopy analysis method. *Appl. Math. Comput.* 2004; **156**, 527-39.
- [25] JH He. Homotopy perturbation method for solving boundary value problems. *Phys. Lett. A* 2006; 350, 87-8.
- [26] A Yildirim and H Kocak. Homotopy perturbation method for solving the space time fractional advection dispersion equation. *Adv. Water. Resour.* 2009; **32**, 1711-6.
- [27] A Yildirim and Y Gulkanat. Analytical approach to fractional Zakharov-kuznetsov equations by He's homotopy perturbation method. *Commun. Theor. Phys.* 2010; 53, 1005-10.

- [28] NA Khan, NU Khan, M Ayaz and A Mahmood. Analytical methods for solving the time fractional Swift Hohenberg (S-H) equation. *Comput. Math. Appl.* 2011; **61**, 2182-5.
- [29] S Kumar and OP Singh. Numerical inversion of Abel integral equation using homotopy perturbation method. *Z. Naturforsch* 2010; **65a**, 677-82.
- [30] Q Wang. Homotopy perturbation method for Kdv equation. *Appl. Math. Comput.* 2007; **90**, 1795-802.
- [31] KA Gepreel. The homotopy perturbation method applied to the nonlinear fractional Kolmogorov-Petrovskii-Piskunov equations. *Appl. Math. Lett.* 2011; **24**, 1428-34.
- [32] F Mainardi. On the Initial Value Problem for the Fractional Diffusion-Wave Equation. In: S Rionero and T Ruggeeri (eds.). Waves and Stability in Continuous Media, World Scientific, Singapore, 1994, p. 246-51.